



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TNFRSF17 (Human) Recombinant Protein

Catalog Number: P9275

Regulation Status: For research use only (RUO)

Product Description: Human TNFRSF17 recombinant protein expressed in *Escherichia coli*.

Sequence:

AGQCSQNEYFDSLHACIPCQLRCSSNTPPLTCQRYC
NASVTNSVKGTNA

Host: *Escherichia coli*

Theoretical MW (kDa): 5.3

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression system

Purification: chromatographic

Purity: > 98% as determined by (a) RP-HPLC.(b) SDS-PAGE.

Storage Buffer: 1 mg protein lyophilized from a solution containing 20 mM sodium phosphate buffer, pH 7.4, 130 mM NaCl. Reconstitute the lyophilized powder in ddH₂O to 100 ug/mL.

Storage Instruction: Lyophilized protein at room temperature for 3 weeks, should be stored at -20°C. Protein aliquots at 4°C for 2-7 days and should be stored at -20°C to -80°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).
Avoid repeated freeze/thaw cycles.

Entrez GeneID: 608

Gene Symbol: TNFRSF17

Gene Alias: BCM, BCMA, CD269

Gene Summary: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is preferentially expressed in mature B lymphocytes, and may be important for B cell development and autoimmune response. This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation. [provided by RefSeq]