



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

TSLP (Human) Recombinant Protein

Catalog Number: P9282

Regulation Status: For research use only (RUO)

Product Description: Human TSLP partial recombinant protein with His tag in N-terminus expressed in *Escherichia coli*.

Sequence:

```
MKHHHHHHASYDFTNCDFEKIKAAAYLSTISKDLITYMS
GTKSTEFNNTVSCSNRPHCLTEIQSLTFNPTAGCASLA
KEMFAMKTKAALAIWCPGYSETQINATQAMKKRRKRK
VTTNKCLEQVSQLQGLWRRFNRPLLKQQ
```

Host: *Escherichia coli*

Theoretical MW (kDa): 16.2

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression system

Purification: chromatographic

Purity: > 95% as determined by SDS-PAGE.

Storage Buffer: Lyophilized from 1X PBS. Reconstitute the lyophilized powder in ddH₂O to 0.5 mg/mL, and is not sterile! Please filter the product by a sterile filter before use.

Storage Instruction: Lyophilized protein should be stored at -20°C. Protein aliquots at 4°C for 2 weeks. Avoid repeated freeze/thaw cycles.

Entrez GeneID: 85480

Gene Symbol: TSLP

Gene Alias: -

Gene Summary: This gene encodes a hemopoietic cytokine proposed to signal through a heterodimeric

receptor complex composed of the thymic stromal lymphopoietin receptor and the IL-7R alpha chain. It mainly impacts myeloid cells and induces the release of T cell-attracting chemokines from monocytes and enhances the maturation of CD11c(+) dendritic cells. Alternative splicing of this gene results in two transcript variants. [provided by RefSeq]