



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CRLF2 (Human) Recombinant Protein

Catalog Number: P9285

Regulation Status: For research use only (RUO)

Product Description: Human CRLF2 partial recombinant protein with His tag in C-terminus expressed in Baculovirus cells.

Sequence:

ADPQGGAEEGVQIIYFNLETVQVTWNASKYSRTNLT
FHYRFNGDEAYDQCTNYLLQEGHTSGCLLDAEQRDDI
LYFSIRNGTHPVFTASRWMVYYLKPSSPKHVRFSWHQ
DAVTVTCSDLSYGDLLYEVQYRSPFDTEWQSKQENT
CNVTIEGLDAEKCYFWVRVKAMEDVYGPDTYPSDW
SEVTCWQRGEIRDACAETPTPPKPKLSKHHHHHH

Host: Viruses

Theoretical MW (kDa): 25.2

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Preparation Method: Baculovirus expression system

Purification: chromatographic

Purity: > 95% as determined by SDS-PAGE.

Storage Buffer: Solution (1 mg/mL) containing 1X PBS, pH 7.4, 10% glycerol.

Storage Instruction: Store at 4°C for one weeks and should be stored at -20°C to -80°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Avoid repeated freeze/thaw cycles.

Entrez GeneID: 64109

Gene Symbol: CRLF2

Gene Alias: CRL2, CRLF2Y, TSLPR

Gene Summary: Cytokine signals are mediated through specific receptor complexes, the components of which are mostly members of the type I cytokine receptor family. Type I cytokine receptors share conserved structural features in their extracellular domain. Receptor complexes are typically heterodimeric, consisting of alpha chains, which provide ligand specificity, and beta (or gamma) chains, which are required for the formation of high-affinity binding sites and signal transduction.[supplied by OMIM]