



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CD207 (Human) Recombinant Protein

Catalog Number: P9352

Regulation Status: For research use only (RUO)

Product Description: Human CD207 partial recombinant protein with His tag in C-terminus expressed in *Escherichia coli*.

Sequence:

ADPPRFMGTISDVKTNVQLLKGRVDNISTLDSEIKKNS
DGMEAAGVQIQMVNESLGYVRSQFLKLTSTVEKANAQ
IQILTRSWEEVSTLNAQIPELKSDEKASALNTKIRALQ
GSLENMSKLLKRQNDILQVVSQGWKYFKGNFYFSLI
PKTWYSAEQFCVSRNSHLTSVTSESEQEFLYKTAGGL
IYWIGLTKAGMEGDWSWVDDTPFNKVQSARFWIPGE
PNNAGNNEHCGNIKAPSLQAWNDAPCDKTFLFICKRP
YVPSEPHHHHHH

Host: Viruses

Theoretical MW (kDa): 30.9

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression system

Purification: chromatographic

Purity: > 90% as determined by SDS-PAGE.

Storage Buffer: Solution (1 mg/mL) containing 1X PBS, pH 7.4, 10% glycerol.

Storage Instruction: Store at 4°C for one weeks and should be stored at -20°C to -80°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).
Avoid repeated freeze/thaw cycles.

Entrez GeneID: 50489

Gene Symbol: CD207

Gene Alias: CLEC4K

Gene Summary: The protein encoded by this gene is expressed only in Langerhans cells which are immature dendritic cells of the epidermis and mucosa. It is localized in the Birbeck granules, organelles present in the cytoplasm of Langerhans cells and consisting of superimposed and zippered membranes. It is a C-type lectin with mannose binding specificity, and it has been proposed that mannose binding by this protein leads to internalization of antigen into Birbeck granules and providing access to a nonclassical antigen-processing pathway. [provided by RefSeq]