



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL27 (Human) Recombinant Protein

Catalog Number: P9428

Regulation Status: For research use only (RUO)

Product Description: Human CCL27 (Q9Y4X3, 25 a.a. - 112 a.a.) partial recombinant protein expressed in *Escherichia coli*.

Sequence:

MFLLPSTACCTQLYRKPLSDKLLRKVIQVELQEADGD
CHLQAFVLHLAQRSICIHQPQNPSLSQWFEHQERKLHG
TLPKLNFGMLRKMG

Host: *Escherichia coli*

Theoretical MW (kDa): 10.3

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression system

Purity: > 95% by SDS-PAGE

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: In 10mM sodium citrate pH 3.5 (10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to -80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 10850

Gene Symbol: CCL27

Gene Alias: ALP, CTACK, CTAK, ESKINE, ILC, PESKY, SCYA27

Gene Summary: This gene is one of several CC cytokine genes clustered on the p-arm of chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The protein encoded by this gene is chemotactic for skin-associated memory T lymphocytes. This cytokine may also play a role in mediating homing of lymphocytes to cutaneous sites. It specifically binds to chemokine receptor 10 (CCR10). Studies of a similar murine protein indicate that these protein-receptor interactions have a pivotal role in T cell-mediated skin inflammation. [provided by RefSeq]