



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL24 (Human) Recombinant Protein

Catalog Number: P9444

Regulation Status: For research use only (RUO)

Product Description: Human CCL24 (O00175, 27 a.a. - 119 a.a.) partial recombinant protein expressed in *Escherichia coli*.

Sequence:

VVIPSPCCMFFVSKRIPENRVVSYQLSSRSTCLKGGVI
FTTKKGQQFCGDPKQEWVQRYMKNLDAKQKKASPR
ARAVA

Host: *Escherichia coli*

Theoretical MW (kDa): 8.800000000000001

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression system

Purity: > 97.0% by SDS-PAGE

Activity: The Biological activity is 50 - 100 ng/mL, was determined by the chemoattract human PBE (peripheral blood eosinophils), corresponding to a specific activity of 10000 - 20000 IU/mg.

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from sterile distilled Water is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to -80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6369

Gene Symbol: CCL24

Gene Alias: Ckb-6, MPIF-2, MPIF2, SCYA24

Gene Summary: This gene belongs to the subfamily of small cytokine CC genes. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity on resting T lymphocytes, a minimal activity on neutrophils, and is negative on monocytes and activated T lymphocytes. The protein is also a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line. [provided by RefSeq]