



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL26 (Human) Recombinant Protein

Catalog Number: P9447

Regulation Status: For research use only (RUO)

Product Description: Human CCL26 (Q9Y258, 24 a.a. - 94 a.a.) partial recombinant protein expressed in *Escherichia coli*.

Sequence:

TRGSDISKTCFQYSHKPLPWTWVRSYEFTSNCSQ
RAVIFTTKRGKVKCTHPRKKWVQKYISLLKTPKQL

Host: *Escherichia coli*

Theoretical MW (kDa): 8.4

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression system

Purity: > 97.0% by SDS-PAGE

Activity: The ED₅₀ is 0.1 - 1.0 ug/mL was determined by the ability to chemoattract BaF3 mouse pro-B cells transfected with mouse CCR3.

Recommend Usage: Biological Activity

SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from sterile distilled Water is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week.

For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 10344

Gene Symbol: CCL26

Gene Alias: IMAC, MGC126714, MIP-4a, MIP-4alpha, SCYA26, TSC-1

Gene Summary: This gene is one of two Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 7. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for normal peripheral blood eosinophils and basophils. The product of this gene is one of three related chemokines that specifically activate chemokine receptor CCR3. This chemokine may contribute to the eosinophil accumulation in atopic diseases. [provided by RefSeq]