



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL1 (Human) Recombinant Protein

Catalog Number: P9479

Regulation Status: For research use only (RUO)

Product Description: Human CCL1 (P22362, 24 a.a. - 96 a.a.) partial recombinant protein with His tag at N-terminus expressed in *Escherichia coli*.

Sequence:

MGSSHHHHHSSGLVPRGSHMKSMQVPFSRCCFSF
AEQEIPLRAILCYRNTSSICSNEGLIFKLKRGKEACALD
TVGWVQRHRKMLRHCPKSRK

Host: *Escherichia coli*

Theoretical MW (kDa): 10.8

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: *Escherichia coli* expression
system

Purity: > 85.0% by SDS-PAGE

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by
the end user.

Storage Buffer: In 20mM Tris-HCl pH 7.5 (1 M DTT, 50
mM NaCl and 10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6346

Gene Symbol: CCL1

Gene Alias: I-309, P500, SCYA1, SISE, TCA3

Gene Summary: This gene is one of several cytokine

genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine is secreted by activated T cells and displays chemotactic activity for monocytes but not for neutrophils. It binds to the chemokine receptor CCR8. [provided by RefSeq]