



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL3 (Human) Recombinant Protein

Catalog Number: P9534

Regulation Status: For research use only (RUO)

Product Description: Human CCL3 (P10147, 23 a.a. - 92 a.a.) partial recombinant protein expressed in *Escherichia coli*.

Sequence:

ASLAADTPTACCFSYTSRQIPQNFADYFETSSQCSKP
GVIFLTKRSRQVCADPSEEWVQKYVSDLELSA

Host: *Escherichia coli*

Theoretical MW (kDa): 7.8

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression system

Purity: > 98.0% by SDS-PAGE

Activity: The Biological activity is 1 - 10 ng/mL, was calculated by the ability of chemo-attraction of Human monocytes, corresponding to a specific activity of 100000 - 1000000 IU/mg.

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from sterile distilled Water
is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to -80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6348

Gene Symbol: CCL3

Gene Alias: G0S19-1, LD78ALPHA, MIP-1-alpha, MIP1A, SCYA3

Gene Summary: Macrophage inflammatory protein-1 is a so-called monokine that is involved in the acute inflammatory state in the recruitment and activation of polymorphonuclear leukocytes (Wolpe et al., 1988 [PubMed 3279154]). Sherry et al. (1988) [PubMed 3058856] demonstrated 2 protein components of MIP1, called by them alpha and beta.[supplied by OMIM]