



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

PF4 (Human) Recombinant Protein

Catalog Number: P9559

Regulation Status: For research use only (RUO)

Product Description: Human PF4 (P02776, 32 a.a. - 101 a.a.) partial recombinant protein with His tag at N-terminus expressed in *Escherichia coli*.

Sequence:

MHHHHHHEAEEDGDLQCLCVKTTTSQVRPRHITSLEVI
KAGPHCPTAQLIATLKNRGRKICLDLQALLYKKIIEHLES

Host: *Escherichia coli*

Theoretical MW (kDa): 8.699999999999999

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression
system

Purity: > 95.0% by SDS-PAGE

Recommend Usage: Biological Activity
SDS-PAGE
The optimal working dilution should be determined by
the end user.

Storage Buffer: Lyophilized from sterile distilled Water
is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 5196

Gene Symbol: PF4

Gene Alias: CXCL4, MGC138298, SCYB4

Gene Summary: Platelet factor-4 is a 70-amino acid
protein that is released from the alpha-granules of

activated platelets and binds with high affinity to heparin. Its major physiologic role appears to be neutralization of heparin-like molecules on the endothelial surface of blood vessels, thereby inhibiting local antithrombin III activity and promoting coagulation. As a strong chemoattractant for neutrophils and fibroblasts, PF4 probably has a role in inflammation and wound repair (Eisman et al., 1990 [PubMed 1695112]).[supplied by OMIM]