



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### PF4 (Human) Recombinant Protein

**Catalog Number:** P9560

**Regulation Status:** For research use only (RUO)

**Product Description:** Human PF4 (P02776, 32 a.a. - 101 a.a.) partial recombinant protein with His tag at N-terminus expressed in *Escherichia coli*.

**Sequence:**

MGSSHHHHHSSGLVPRGSHMEAEEDGDLQCLCVKT  
TSQVRPRHITSLEVIKAGPHCPTAQLIATLKNRKCILD  
LQAPLYKKIIKKLLES

**Host:** *Escherichia coli*

**Theoretical MW (kDa):** 10

**Protocols:** See our web site at  
<http://www.abnova.com/support/protocols.asp> or product  
page for detailed protocols

**Form:** Liquid

**Preparation Method:** *Escherichia coli* expression  
system

**Purity:** > 85.0% by SDS-PAGE

**Recommend Usage:** Biological Activity  
SDS-PAGE  
The optimal working dilution should be determined by  
the end user.

**Storage Buffer:** In 20mM Tris-HCl pH 8.0 (0.2 M NaCl,  
2 mM DTT and 50% glycerol)

**Storage Instruction:** Store at 2°C to 8°C for 1 week.  
For long term storage, aliquot and store at -20°C to  
-80°C.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 5196

**Gene Symbol:** PF4

**Gene Alias:** CXCL4, MGC138298, SCYB4

**Gene Summary:** Platelet factor-4 is a 70-amino acid

protein that is released from the alpha-granules of activated platelets and binds with high affinity to heparin. Its major physiologic role appears to be neutralization of heparin-like molecules on the endothelial surface of blood vessels, thereby inhibiting local antithrombin III activity and promoting coagulation. As a strong chemoattractant for neutrophils and fibroblasts, PF4 probably has a role in inflammation and wound repair (Eisman et al., 1990 [PubMed 1695112]).[supplied by OMIM]