



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCL17 (Human) Recombinant Protein

Catalog Number: P9582

Regulation Status: For research use only (RUO)

Product Description: Human CCL17 (Q92583, 24 a.a. - 94 a.a.) partial recombinant protein expressed in *Escherichia coli*.

Sequence:

ARGTNVGRECCLEYFKGAIPLRKLKTWYQTSSEDCSRD
AIVFVTVQGRAICSDPNNKRVKNAVLYLQSLERS

Host: *Escherichia coli*

Theoretical MW (kDa): 8

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: *Escherichia coli* expression system

Purity: > 97.0% by SDS-PAGE

Activity: The Biological activity is 1.0 - 10 ng/mL, was determined by the ability to chemoattract human T-Lymphocytes, corresponding to a specific activity of 100000 - 1000000 IU/mg.

Recommend Usage: Biological Activity
SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from sterile distilled Water is 0.5 mg/mL

Storage Instruction: Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 6361

Gene Symbol: CCL17

Gene Alias: A-152E5.3, ABCD-2, MGC138271, MGC138273, SCYA17, TARC

Gene Summary: This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for T lymphocytes, but not monocytes or granulocytes. The product of this gene binds to chemokine receptors CCR4 and CCR8. This chemokine plays important roles in T cell development in thymus as well as in trafficking and activation of mature T cells. [provided by RefSeq]