



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

FCER2 (Human) Recombinant Protein

Catalog Number: P9662

Regulation Status: For research use only (RUO)

Product Description: Human FCER2 (P06734, 48 a.a. - 321 a.a.) partial recombinant protein with His tag at C-terminus expressed in Sf9 cells.

Sequence:

ADPDTTQSLKQLEERAARNVSQVSKNLESHHGQMA
KKSQSTQISQELEELRAEQRLKSQDLELSWNLNGLQ
ADLSSFKSQELNERNEASDLLERLREEVTKLRMELQV
SSGFVCNTCPEKWINFQRKCYFYGKGTKQWVHARYA
CDDMEGQLVSIHSPEEQDFLTKHASHTGSWIGLR NLD
LKGEFIWVDGSHVDYSNWAPGEPTSRSQGEDCVMM
RSGRWNDAFCDRKLGAWVCDRLATCTPPASEGSA
ESMGPDSPDPDGRLLPT PSAPLHSHHHHHH

Host: insect

Theoretical MW (kDa): 32

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product
page for detailed protocols

Form: Liquid

Preparation Method: Sf9 cell expression system

Purity: > 90.0% by SDS-PAGE

Recommend Usage: Biological Activity
SDS-PAGE
The optimal working dilution should be determined by
the end user.

Storage Buffer: In PBS pH 7.4 (10% glycerol)

Storage Instruction: Store at 2°C to 8°C for 2-4 week.
For long term storage, aliquot and store at -20°C to
-80°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 2208

Gene Symbol: FCER2

Gene Alias: CD23, CD23A, CLEC4J, FCE2, IGEBF

Gene Summary: The human leukocyte differentiation antigen CD23 (FCE2) is a key molecule for B-cell activation and growth. It is the low-affinity receptor for IgE. The truncated molecule can be secreted, then functioning as a potent mitogenic growth factor.[supplied by OMIM]