



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### CD300C (Human) Recombinant Protein

**Catalog Number:** P9686

**Regulation Status:** For research use only (RUO)

**Product Description:** Human CD300C (Q08708, 21 a.a. - 183 a.a.) partial recombinant protein with His tag at C-terminus expressed in Sf9 cells.

**Sequence:**

```
MTVAGPVGGSLSVQCRYEKEHRTLNKFWCRPPQILR  
CDKIVETKGSAGKRNGRVSIRDSPANLSFTVTLENLTE  
EDAGTYWCGVDTPWLRDFHDPIVEVEVSVFPAGTTTA  
SSPQSSMGTS GPPTKLPVHTWPSVTRKDSPEPSPHP  
GSLFSNVRLEHHHHHHH
```

**Host:** insect

**Theoretical MW (kDa):** 18

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Preparation Method:** Sf9 cell expression system

**Purity:** > 95.0% by SDS-PAGE

**Recommend Usage:** Biological Activity

SDS-PAGE

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS pH 7.4 (10% glycerol)

**Storage Instruction:** Store at 2°C to 8°C for 2-4 week.

For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 10871

**Gene Symbol:** CD300C

**Gene Alias:** CMRF-35A, CMRF35, CMRF35A, CMRF35A1, IGSF16, LIR

**Gene Summary:** The CMRF35 antigen, which was identified by reactivity with a monoclonal antibody, is present on monocytes, neutrophils, and some T and B lymphocytes (Jackson et al., 1992 [PubMed 1349532]).[supplied by OMIM]