

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

#### **Datasheet**

#### **CD44 (Human) Recombinant Protein**

Catalog Number: P9696

Regulation Status: For research use only (RUO)

**Product Description:** Human CD44 (P16070-1, Gln 21 - Pro 220 & Gln 386 - Ala 427) partial recombinant protein with His tag at C-terminus expressed in HEK293 cells.

Sequence: Gln 21-Pro 220 & Gln 386-Ala 427

Host: Human

Theoretical MW (kDa): 27.91

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Liquid

**Preparation Method:** Mammalian cell (Expi293, highyield transient HEK293) expression system

Purity: > 95% as determined by Tris-Bis PAGE; > 95%

as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein

(determined by LAL method)

Activity: The EC<sub>50</sub> was 11.8 ng/mL, messured by ELISA

at 0.5 ug/mL.

Recommend Usage: Biological Activity

ELISA SDS-PAGE

The optimal working dilution should be determined by

the end user.

**Storage Buffer:** In PBS pH 7.4

**Storage Instruction:** Store at -80°C for 12 Month. Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 960

Gene Symbol: CD44

Gene Alias: CDW44, CSPG8, ECMR-III, HCELL, IN, LHR, MC56, MDU2, MDU3, MGC10468, MIC4, MUTCH-I, Pgp1

Gene Summary: The protein encoded by this gene is a cell-surface glycoprotein involved in cell-cell interactions, cell adhesion and migration. It is a receptor for hyaluronic acid (HA) and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). This protein participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, hematopoiesis, and tumor metastasis. Transcripts for this gene undergo complex alternative splicing that results in many functionally distinct isoforms, however, the full length nature of some of these variants has not been determined. Alternative splicing is the basis for the structural and functional diversity of this protein, and may be related to tumor metastasis. [provided by RefSeq]