



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCR8 (Human) Recombinant Protein

Catalog Number: P9721

Regulation Status: For research use only (RUO)

Product Description: Human CCR8 (P51685-1, 1 a.a. - 35 a.a.) partial recombinant protein with hFc tag at C-terminus expressed in HEK293 cells.

Sequence:

MDYTLDL SVTTVTDY YYPDIFSSPCDAELIQTNGK

Host: Human

Theoretical MW (kDa): 30.62

Protocols: See our web site at

<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (Expi293, high-yield transient HEK293) expression system

Purity: > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein (determined by LAL method)

Activity: The EC₅₀ was 17.6 ng/mL, measured by ELISA at 1 ug/mL.

Recommend Usage: Biological Activity

ELISA

SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from sterile distilled Water is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week.

For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 1237

Gene Symbol: CCR8

Gene Alias: CDw198, CKR-L1, CKRL1, CMKBR8, CMKBRL2, CY6, GPR-CY6, MGC129966, MGC129973, TER1

Gene Summary: This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptors are important for the migration of various cell types into the inflammatory sites. This receptor protein preferentially expresses in the thymus. I-309, thymus activation-regulated cytokine (TARC) and macrophage inflammatory protein-1 beta (MIP-1 beta) have been identified as ligands of this receptor. Studies of this receptor and its ligands suggested its role in regulation of monocyte chemotaxis and thymic cell apoptosis. More specifically, this receptor may contribute to the proper positioning of activated T cells within the antigenic challenge sites and specialized areas of lymphoid tissues. This gene is located at the chemokine receptor gene cluster region. [provided by RefSeq]