



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### IL11RA (Human) Recombinant Protein

**Catalog Number:** P9765

**Regulation Status:** For research use only (RUO)

**Product Description:** Human IL11RA (Q14626-1, Ser24-Ala370) partial recombinant protein with His tag at C-Terminus expressed in HEK293 cells.

**Sequence:** Ser24-Ala370

**Host:** Human

**Theoretical MW (kDa):** 38.8

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Preparation Method:** Mammalian cell (HEK293) expression system

**Recommend Usage:** Biological Activity  
SPR  
SDS-PAGE  
The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS pH 7.4 (8% trehalose)

**Storage Instruction:** Store at -80°C for 12 Month.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 3590

**Gene Symbol:** IL11RA

**Gene Alias:** MGC2146

**Gene Summary:** Interleukin 11 is a stromal cell-derived cytokine that belongs to a family of pleiotropic and redundant cytokines that use the gp130 transducing subunit in their high affinity receptors. This gene encodes the IL-11 receptor, which is a member of the hematopoietic cytokine receptor family. This particular receptor is very similar to ciliary neurotrophic factor,

since both contain an extracellular region with a 2-domain structure composed of an immunoglobulin-like domain and a cytokine receptor-like domain. Alternative splicing has been observed at this locus, and three variants encoding two different isoforms have been identified. [provided by RefSeq]