



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### IL17RA (Human) Recombinant Protein

**Catalog Number:** P9776

**Regulation Status:** For research use only (RUO)

**Product Description:** Human IL17RA (Q96F46-1, Leu33-Trp320) partial recombinant protein with His tag at C-Terminus expressed in HEK293 cells.

**Sequence:** Leu33-Trp320

**Host:** Human

**Theoretical MW (kDa):** 34.6

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Lyophilized

**Preparation Method:** Mammalian cell (HEK293) expression system

**Recommend Usage:** Biological Activity

ELISA

SPR

SDS-PAGE

The optimal working dilution should be determined by the end user.

**Storage Buffer:** Lyophilized from sterile distilled Water is > 100 ug/mL

**Storage Instruction:** Store at 2°C to 8°C for 1 week.

For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 23765

**Gene Symbol:** IL17RA

**Gene Alias:** CD217, CDw217, IL-17RA, IL17R, MGC10262, hIL-17R

**Gene Summary:** Interleukin 17A (IL17A) is a proinflammatory cytokine secreted by activated T-

lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure. [provided by RefSeq]