

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





9F, No. 108, Jhouzih St.,Taipei, Taiwan Tel: + 886-2-8751-1888 Fax: + 886-2-6602-1218 E-mail: sales@abnova.com

#### **Datasheet**

#### **DDR1 (Human) Recombinant Protein**

Catalog Number: P9802

Regulation Status: For research use only (RUO)

**Product Description:** Human DDR1(Q08345-1, Asp21-Ala417) partial recombinant protein with His tag

at C-terminus expressed in HEK293 cells.

Sequence: Asp21-Ala417

Host: Human

Theoretical MW (kDa): 45.1

Protocols: See our web site at

http://www.abnova.com/support/protocols.asp or product

page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293)

expression system

**Purity:** > 95% as determined by Tris-Bis PAGE; > 95%

as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein

(determined by LAL method)

**Activity:** The EC<sub>50</sub> was 5.9 ng/mL, messured by ELISA at 0.5 ug/mL. (QC Test). The affinity constant of 0.806 nM as determined in SPR assay (Biacore T200)

Recommend Usage: Biological Activity

ELISA SDS-PAGE SPR

The optimal working dilution should be determined by

the end user.

Storage Buffer: Lyophilized from sterile distilled Water

is > 100 ug/mL

**Storage Instruction:** Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to

-80°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GenelD: 780

Gene Symbol: DDR1

Gene Alias: CAK, CD167, DDR, EDDR1, MCK10, NEP,

NTRK4, PTK3, PTK3A, RTK6, TRKE

Gene Summary: Receptor tyrosine kinases (RTKs) play a key role in the communication of cells with their microenvironment. These molecules are involved in the regulation of cell growth, differentiation and metabolism. The protein encoded by this gene is a RTK that is widely expressed in normal and transformed epithelial cells and is activated by various types of collagen. This protein belongs to a subfamily of tyrosine kinase receptors with a homology region to the Dictyostelium discoideum protein discoidin I in their extracellular domain. Its autophosphorylation is achieved by all collagens so far tested (type I to type VI). In situ studies and Northernblot analysis showed that expression of this encoded protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, this protein is significantly over-expressed in several human tumors from breast, ovarian, esophageal, and pediatric brain. This gene is located on chromosome 6p21.3 in proximity to several HLA class I genes. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq]