



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### CD8A (Human) Recombinant Protein

**Catalog Number:** P9829

**Regulation Status:** For research use only (RUO)

**Product Description:** Human CD8A (P01732-1, Ser22-Asp182) partial recombinant protein with His-Avi tag at C-Terminus expressed in HEK293 cells.

**Sequence:** Ser22-Asp182

**Host:** Human

**Theoretical MW (kDa):** 20.5

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Lyophilized

**Preparation Method:** Mammalian cell (HEK293) expression system

**Purity:** > 95% as determined by Tris-Bis PAGE

**Endotoxin Level:** < 1 EU per 1 ug of protein (determined by LAL method)

**Activity:** The EC<sub>50</sub> was 4.8 ng/mL, measured by ELISA at 0.5 ug/mL.

**Recommend Usage:** Biological Activity

ELISA

SDS-PAGE

The optimal working dilution should be determined by the end user.

**Storage Buffer:** Lyophilized from sterile distilled Water is > 100 ug/mL

**Storage Instruction:** Store at 2°C to 8°C for 1 week.

For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 925

**Gene Symbol:** CD8A

**Gene Alias:** CD8, Leu2, MAL, p32

**Gene Summary:** The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen acts as a corepressor with the T-cell receptor on the T lymphocyte to recognize antigens displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The coreceptor functions as either a homodimer composed of two alpha chains, or as a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 alpha chain isoforms. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]