



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### BST1 (Human) Recombinant Protein

**Catalog Number:** P9888

**Regulation Status:** For research use only (RUO)

**Product Description:** Human BST1 (Q10588-1, Gly29-Ala293) partial recombinant protein with His tag at C-Terminus expressed in HEK293 cells.

**Sequence:** Gly29-Ala293

**Host:** Human

**Theoretical MW (kDa):** 31.2

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Lyophilized

**Preparation Method:** Mammalian cell (HEK293) expression system

**Purity:** > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

**Endotoxin Level:** < 1 EU per 1 ug of protein (determined by LAL method)

**Activity:** The EC<sub>50</sub> was 8.4 ng/mL, measured by ELISA at 0.5 ug/mL.

**Recommend Usage:** Biological Activity

ELISA

SDS-PAGE

The optimal working dilution should be determined by the end user.

**Storage Buffer:** Lyophilized from sterile distilled Water is > 100 ug/mL

**Storage Instruction:** Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C. Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 683

**Gene Symbol:** BST1

**Gene Alias:** CD157

**Gene Summary:** Bone marrow stromal cell antigen-1 is a stromal cell line-derived glycosylphosphatidylinositol-anchored molecule that facilitates pre-B-cell growth. The deduced amino acid sequence exhibits 33% similarity with CD38. BST1 expression is enhanced in bone marrow stromal cell lines derived from patients with rheumatoid arthritis. The polyclonal B-cell abnormalities in rheumatoid arthritis may be, at least in part, attributed to BST1 overexpression in the stromal cell population. [provided by RefSeq]