



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IL5RA (Human) Recombinant Protein

Catalog Number: P9898

Regulation Status: For research use only (RUO)

Product Description: Human IL5RA (Q01344-1, Asp21-Glu335) partial recombinant protein with hFc tag at C-Terminus expressed in HEK293 cells.

Sequence: Asp21-Glu335

Host: Human

Theoretical MW (kDa): 62.6

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein (determined by LAL method)

Activity: The EC₅₀ was 6.9 ng/mL, measured by ELISA at 0.5 ug/mL.

Recommend Usage: Biological Activity

ELISA

SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from sterile distilled Water is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week.

For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3568

Gene Symbol: IL5RA

Gene Alias: CD125, CDw125, HSIL5R3, IL5R, MGC26560

Gene Summary: The protein encoded by this gene is an interleukin 5 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL5 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL5. This protein has been found to interact with syndecan binding protein (syntenin), which is required for IL5 mediated activation of the transcription factor SOX4. Six alternatively spliced transcript variants encoding three distinct isoforms have been reported. [provided by RefSeq]