



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

OSMR (Human) Recombinant Protein

Catalog Number: P9939

Regulation Status: For research use only (RUO)

Product Description: Human OSMR (Q99650-1, Glu28-Met740) partial recombinant protein with His tag at C-Terminus expressed in HEK293 cells.

Sequence: Glu28-Met740

Host: Human

Theoretical MW (kDa): 82.2

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Lyophilized

Preparation Method: Mammalian cell (HEK293) expression system

Purity: > 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC

Endotoxin Level: < 1 EU per 1 ug of protein (determined by LAL method)

Activity: The EC₅₀ was 4.4 ng/mL, measured by ELISA at 1 ug/mL.

Recommend Usage: Biological Activity

ELISA

SDS-PAGE

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from sterile distilled Water is > 100 ug/mL

Storage Instruction: Store at 2°C to 8°C for 1 week. For long term storage, aliquot and store at -20°C to -80°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 9180

Gene Symbol: OSMR

Gene Alias: MGC150626, MGC150627, MGC75127, OSMRB

Gene Summary: Oncostatin M is a member of the IL6 family of cytokines. Functional receptors for IL6 family cytokines are multisubunit complexes involving members of the hematopoietin receptor superfamily. Many IL6 cytokines utilize gp130 as a common receptor subunit. OSM binds to the gp130 receptor subunit and, in association with the leukemia inhibitory factor receptor, induces a proliferative response in permissive cells. OSMR is an alternative subunit (for an OSM receptor complex (a heterodimer of gp130 and OSMR) that is activated by OSM but not by LIF [provided by RefSeq]