



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

Rabbit Fab Anti-Human IgG (H&L) secondary antibody (FITC)

Catalog Number: PAB31782

Regulation Status: For research use only (RUO)

Product Description: Rabbit Fab polyclonal antibody raised against human IgG (H&L). The antibody is conjugated with Fluorescein (FITC).

Immunogen: Human IgG whole molecule.

Host: Rabbit

Reactivity: Human

Applications: FLISA, Flow Cyt, IF
(See our web site product page for detailed applications information)

Protocols: See our web site at
<http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: IgG (H&L). Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-fluorescein and anti-rabbit serum. No reaction was observed against anti-papain or anti-rabbit IgG F(c).

Form: Lyophilized

Conjugation: FITC

Purification: This antibody was prepared from monospecific antiserum by immunoaffinity chromatography using human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation.

Isotype: IgG Fab

Conjugation Note: FP Value: 1.3 moles Fluorescein (FITC) per mole of IgG Fab

Recommend Usage: FLISA (1:10000-1:50000)
Flow Cytometry (1:500-1:2500)
Immunofluorescence (1:1000-1:5000)

The optimal working dilution should be determined by the end user.

Storage Buffer: Lyophilized from 0.02 M potassium phosphate, 0.15 M sodium chloride, pH 7.2 (10 mg/mL BSA (immunoglobulin and protease free), 0.01% (w/v) sodium azide).

Storage Instruction: Store vial at 4°C prior to restoration.

After reconstitution with 500 μ L deionized water (or equivalent), aliquot contents and freeze at -20°C or below for extended storage.

Aliquot to avoid repeated freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. Dilute only prior to immediate use.