



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

EGFR recombinant monoclonal antibody, clone 2F8 (Zalutumumab; HuMax-EGFR)

Catalog Number: RAB01158

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human EGFR.

Clone Name: 2F8

Immunogen: Original antibody is raised against by immunizing HuMAb mice with alternating A431 cells and purified EGFR administration.

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: This antibody binds domain III of the human epidermal growth factor receptor (EGFR).

Form: Liquid

Purification: Protein A affinity purification

Isotype: IgG, Kappa

Recommend Usage: Flow cytometry

Immunofluorescence

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS with 0.02% Proclin 300

Storage Instruction: Store at 4°C for up to 3 months.
For longer storage, aliquot and store at -20°C.

Entrez GeneID: 1956

Gene Symbol: EGFR

Gene Alias: ERBB, ERBB1, HER1, PIG61, mENA

Gene Summary: The protein encoded by this gene is a transmembrane glycoprotein that is a member of the

protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Mutations in this gene are associated with lung cancer. [provided by RefSeq]