



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CASP6 recombinant monoclonal antibody, clone R01-3A0

Catalog Number: RAB01478

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human CASP6.

Clone Name: R01-3A0

Immunogen: Original antibody is raised against recombinant protein corresponding to human CASP6.

Theoretical MW (kDa): Calculated MW: 33 kD

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Affinity purification

Isotype: IgG

Recommend Usage: Immunohistochemistry
(Formalin/PFA-fixed paraffin-embedded sections)
Western Blot

The optimal working dilution should be determined by the end user.

Storage Buffer: In 50mM Tris-Glycine, pH 7.4, (0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction: Store at 4°C. For longer storage, aliquot and store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 839

Gene Symbol: CASP6

Gene Alias: MCH2

Gene Summary: This gene encodes a protein which is

a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein is processed by caspases 7, 8 and 10, and is thought to function as a downstream enzyme in the caspase activation cascade. Alternative splicing of this gene results in two transcript variants that encode different isoforms. [provided by RefSeq]