



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

NCOA1 recombinant monoclonal antibody, clone R07-5B7

bHLHe42, bHLHe74

Catalog Number: RAB01713**Regulatory Status:** For research use only (RUO)**Product Description:** Rabbit recombinant monoclonal antibody raised against human NCOA1.**Clone Name:** R07-5B7**Immunogen:** Original antibody is raised against recombinant protein corresponding to human SRC1**Theoretical MW (kDa):** Calculated MW: 157 k**Antibody Species:** Rabbit**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols**Form:** Liquid**Purification:** Affinity purification**Isotype:** IgG**Recommend Usage:** Immunohistochemistry

(1:50-1:100)

Immunoprecipitation (1:20)

Western Blot (1:500-1:1,000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)**Storage Instruction:** Store at 4°C for short term. For long term storage store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 8648**Gene Symbol:** NCOA1**Gene Alias:** F-SRC-1, KAT13A, MGC129719, MGC129720, NCoA-1, RIP160, SRC-1, SRC1,**Gene Summary:** The protein encoded by this gene acts as a transcriptional coactivator for steroid and nuclear hormone receptors. It is a member of the p160/steroid receptor coactivator (SRC) family and like other family members has histone acetyltransferase activity and contains a nuclear localization signal, as well as bHLH and PAS domains. The product of this gene binds nuclear receptors directly and stimulates the transcriptional activities in a hormone-dependent fashion. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq]