



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### RELA recombinant monoclonal antibody, clone R09-3H2

**Catalog Number:** RAB02061

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human RELA.

**Clone Name:** R09-3H2

**Immunogen:** Original antibody is raised against a synthetic peptide corresponding to human RELA.

**Theoretical MW (kDa):** Calculated MW: 60 kD

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Affinity purification

**Isotype:** IgG

**Recommend Usage:** Immunofluorescence (1:50-1:200)  
Immunohistochemistry (1:50-1:100)  
Immunoprecipitation (1:20)  
Western Blot (1:500-1:1000)  
The optimal working dilution should be determined by the end user.

**Storage Buffer:** In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

**Storage Instruction:** Store at -20 °C.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 5970

**Gene Symbol:** RELA

**Gene Alias:** MGC131774, NFKB3, p65

**Gene Summary:** NFKB1 (MIM 164011) or NFKB2 (MIM

164012) is bound to REL (MIM 164910), RELA, or RELB (MIM 604758) to form the NFκB complex. The p50 (NFKB1)/p65 (RELA) heterodimer is the most abundant form of NFκB. The NFκB complex is inhibited by I-κB proteins (NFKBIA, MIM 164008 or NFKBIB, MIM 604495), which inactivate NFκB by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-κB proteins by kinases (IKBKA, MIM 600664, or IKBKB, MIM 603258) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NFκB complex. Activated NFκB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).[supplied by OMIM]