



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### LCK recombinant monoclonal antibody, clone R01-8A5

**Catalog Number:** RAB02136

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human LCK.

**Clone Name:** R01-8A5

**Immunogen:** Original antibody is raised against recombinant protein corresponding to human LCK.

**Theoretical MW (kDa):** Calculated MW: 58 kD

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Affinity purification

**Isotype:** IgG

**Recommend Usage:** Immunocytochemistry

Immunofluorescence(1:50-1:200)

Immunohistochemistry (1:50-1:100)

Immunoprecipitation(1:20)

Western Blot (1:500-1:1000)

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

**Storage Instruction:** Store at -20 °C.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 3932

**Gene Symbol:** LCK

**Gene Alias:** YT16, p56lck, pp58lck

**Gene Summary:** This gene is a member of the Src family of protein tyrosine kinases (PTKs). The encoded protein is a key signaling molecule in the selection and maturation of developing T-cells. It contains N-terminal sites for myristylation and palmitoylation, a PTK domain, and SH2 and SH3 domains which are involved in mediating protein-protein interactions with phosphotyrosine-containing and proline-rich motifs, respectively. The protein localizes to the plasma membrane and pericentrosomal vesicles, and binds to cell surface receptors, including CD4 and CD8, and other signaling molecules. Multiple alternatively spliced variants, encoding the same protein, have been described. [provided by RefSeq]