



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

IKBKB recombinant monoclonal antibody, clone R01-9H6

Catalog Number: RAB02178

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human IKBKB.

Clone Name: R01-9H6

Immunogen: Original antibody is raised against a synthetic peptide corresponding to human IKBKB.

Theoretical MW (kDa): Calculated MW: 87 kD

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Affinity purification

Isotype: IgG

Recommend Usage: Immunofluorescence(1:50-1:200)
Western Blot (1:500-1:1000)
The optimal working dilution should be determined by the end user.

Storage Buffer: In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction: Store at -20 °C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3551

Gene Symbol: IKBKB

Gene Alias: FLJ40509, IKK-beta, IKK2, IKKB, MGC131801, NFKBIKB

Gene Summary: NFKB1 (MIM 164011) or NFKB2 (MIM 164012) is bound to REL (MIM 164910), RELA (MIM

164014), or RELB (MIM 604758) to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA, MIM 164008, or NFKBIB, MIM 604495), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, MIM 600664, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).[supplied by OMIM]