



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### CYCS recombinant monoclonal antibody, clone R02-9A7

**Catalog Number:** RAB02342

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human CYCS.

**Clone Name:** R02-9A7

**Immunogen:** Original antibody is raised against a synthetic peptide corresponding to human CYCS.

**Theoretical MW (kDa):** Calculated MW: 12 kD

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Affinity purification

**Isotype:** IgG

**Recommend Usage:** Immunocytochemistry

Immunofluorescence (1:50-1:200)

Immunoprecipitation (1:20)

Western Blot (1:500-1:1000)

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In 50 mM Tris-Glycine, pH 7.4 (0.15 M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA)

**Storage Instruction:** Store at -20 °C.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 54205

**Gene Symbol:** CYCS

**Gene Alias:** CYC, HCS

**Gene Summary:** This gene encodes cytochrome c, a

component of the electron transport chain in mitochondria. The heme group of cytochrome c accepts electrons from the b-c1 complex and transfers electrons to the cytochrome oxidase complex. Cytochrome c is also involved in initiation of apoptosis. Upon release of cytochrome c to the cytoplasm, the protein binds apoptotic protease activating factor which activates the apoptotic initiator procaspase 9. Many cytochrome c pseudogenes exist, scattered throughout the human genome. [provided by RefSeq]