



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

Active Caspase-3 recombinant monoclonal antibody

Catalog Number: RAB02394

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against recombinant Active Caspase-3.

Theoretical MW (kDa): 17

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: This antibody detects endogenous levels of Activated-Caspase-3 p17 protein.

Form: Liquid

Purification: Protein A purification

Isotype: IgG

Recommend Usage: Immunocytochemistry (1:100-1:300)
Immunofluorescence (1:100-1:300)
Immunohistochemistry (1:50-1:200)
Western Blot (1:500-1:2000)

Storage Buffer: In PBS, pH 7.2 (0.02% sodium azide and 50% glycerol)

Storage Instruction: Store at 4°C short term.
Aliquot and store at -20°C long term.
Avoid freeze-thaw cycles.

Entrez GeneID: 836

Gene Symbol: CASP3

Gene Alias: CPP32, CPP32B, SCA-1

Gene Summary: This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell

apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 6, 7 and 9, and the protein itself is processed by caspases 8, 9 and 10. It is the predominant caspase involved in the cleavage of amyloid-beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease. Alternative splicing of this gene results in two transcript variants that encode the same protein. [provided by RefSeq]