



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CD86 recombinant monoclonal antibody

Catalog Number: RAB02398

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against recombinant CD86.

Theoretical MW (kDa): 37, 70

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: This antibody detects endogenous levels of CD86 and does not cross-react with related proteins.

Form: Liquid

Purification: Protein A purification

Isotype: IgG

Recommend Usage: Immunocytochemistry

(1:50-1:200)

Immunofluorescence (1:50-1:200)

Immunohistochemistry (1:50-1:200)

Western Blot (1:500-1:2000)

Storage Buffer: In PBS, pH 7.2 (0.02% sodium azide and 50% glycerol)

Storage Instruction: Store at 4°C short term.

Aliquot and store at -20°C long term.

Avoid freeze-thaw cycles.

Entrez GeneID: 942

Gene Symbol: CD86

Gene Alias: B7-2, B70, CD28LG2, LAB72, MGC34413

Gene Summary: This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at

the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in two transcript variants encoding different isoforms. Additional transcript variants have been described, but their full-length sequences have not been determined. [provided by RefSeq]