



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

BCL2 recombinant monoclonal antibody

Catalog Number: RAB02405

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against recombinant BCL2.

Immunogen: Original antibody is raised against a synthetic peptide of human Bcl-2

Theoretical MW (kDa): 27

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: Recognizes endogenous levels of BCL2 protein.

Form: Liquid

Purification: Immunogen affinity chromatography

Isotype: IgG

Recommend Usage: Immunohistochemistry (1:50-1:100)
Immunoprecipitation(1:10-1:50)
Western Blot (1:500-1:1000)

Storage Buffer: In 50mM Tris-Glycine, pH 7.4 (0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction: Store at 4°C short term.
Aliquot and store at -20°C long term.
Avoid freeze-thaw cycles.

Entrez GeneID: 596

Gene Symbol: BCL2

Gene Alias: Bcl-2

Gene Summary: This gene encodes an integral outer

mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends. [provided by RefSeq]