



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

EIF4EBP1 (Phospho T46) recombinant monoclonal antibody

Catalog Number: RAB02454

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human EIF4EBP1.

Immunogen: Original antibody is raised against a synthetic phosphopeptide corresponding to residues surrounding Thr45 of human eIF4EBP1.

Theoretical MW (kDa): 15

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: Recognizes endogenous levels of 4EBP1 (pT46) protein.

Form: Liquid

Purification: Immunogen affinity chromatography

Isotype: IgG

Recommend Usage: Immunohistochemistry

(1:50-1:100)

Immunoprecipitation(1:10-1:50)

Western Blot (1:500-1:1000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In 50mM Tris-Glycine, pH 7.4 (0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction: Store at 4°C short term.

Aliquot and store at -20°C long term.

Avoid freeze-thaw cycles.

Entrez GeneID: 1978

Gene Symbol: EIF4EBP1

Gene Alias: 4E-BP1, 4EBP1, BP-1, MGC4316, PHAS-I

Gene Summary: This gene encodes one member of a family of translation repressor proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation. [provided by RefSeq]