



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

UPF1 recombinant monoclonal antibody

Catalog Number: RAB02536

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human UPF1.

Immunogen: Original antibody is raised against a synthetic peptide of human UPF1.

Theoretical MW (kDa): 120

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: Recognizes endogenous levels of UPF1 protein.

Form: Liquid

Purification: Immunogen affinity chromatography

Isotype: IgG

Recommend Usage: Immunocytochemistry (1:50-1:100)

Immunofluorescence (1:50-1:100)

Immunohistochemistry (1:50-1:100)

Immunoprecipitation (1:10-1:50)

Western Blot (1:500-1:1000)

Storage Buffer: In 50mM Tris-Glycine, pH 7.4 (0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA)

Storage Instruction: Store at 4°C short term.
Aliquot and store at -20°C long term.
Avoid freeze-thaw cycles.

Entrez GeneID: 5976

Gene Symbol: UPF1

Gene Alias: FLJ43809, FLJ46894, HUPF1, KIAA0221,

NORF1, RENT1, pNORF1

Gene Summary: This gene encodes a protein that is part of a post-splicing multiprotein complex involved in both mRNA nuclear export and mRNA surveillance. mRNA surveillance detects exported mRNAs with truncated open reading frames and initiates nonsense-mediated mRNA decay (NMD). When translation ends upstream from the last exon-exon junction, this triggers NMD to degrade mRNAs containing premature stop codons. This protein is located only in the cytoplasm. When translation ends, it interacts with the protein that is a functional homolog of yeast Upf2p to trigger mRNA decapping. Use of multiple polyadenylation sites has been noted for this gene. [provided by RefSeq]