



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### IMPDH2 recombinant monoclonal antibody

**Catalog Number:** RAB02594

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human IMPDH2.

**Immunogen:** Original antibody is raised against a synthetic peptide of human IMPDH2.

**Theoretical MW (kDa):** 55

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Specificity:** Recognizes endogenous levels of IMPDH2 protein.

**Form:** Liquid

**Purification:** Immunogen affinity chromatography

**Isotype:** IgG

**Recommend Usage:** Immunocytochemistry (1:50-1:100)

Immunofluorescence (1:50-1:100)

Immunohistochemistry (1:50-1:100)

Immunoprecipitation (1:10-1:50)

Western Blot (1:500-1:1000)

**Storage Buffer:** In 50mM Tris-Glycine, pH 7.4 (0.15M NaCl, 50% Glycerol, 0.01% Sodium azide and 0.05% BSA)

**Storage Instruction:** Store at 4°C short term.  
Aliquot and store at -20°C long term.  
Avoid freeze-thaw cycles.

**Entrez GeneID:** 3615

**Gene Symbol:** IMPDH2

**Gene Alias:** IMPD2, IMPDH-II

**Gene Summary:** This gene encodes the rate-limiting enzyme in the de novo guanine nucleotide biosynthesis. It is thus involved in maintaining cellular guanine deoxy- and ribonucleotide pools needed for DNA and RNA synthesis. The encoded protein catalyzes the NAD-dependent oxidation of inosine-5'-monophosphate into xanthine-5'-monophosphate, which is then converted into guanosine-5'-monophosphate. This gene is up-regulated in some neoplasms, suggesting it may play a role in malignant transformation. [provided by RefSeq]