



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CCNE2 recombinant monoclonal antibody

Catalog Number: RAB02631

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human CCNE2.

Immunogen: Original antibody is raised against recombinant human Cyclin E2 protein, around C-terminal 100aa.

Theoretical MW (kDa): 44

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: This antibody detects endogenous levels of Cyclin E2 protein.

Form: Liquid

Purification: Protein A purification

Isotype: IgG

Recommend Usage: Flow Cytometry (2 ug)
The optimal working dilution should be determined by the end user.

Storage Buffer: In 0.01M TBS, pH7.4 (1% BSA, 0.03% Proclin300 and 50% Glycerol)

Storage Instruction: Store at 4°C short term.
Aliquot and store at -20°C long term.
Avoid freeze-thaw cycles.

Entrez GeneID: 9134

Gene Symbol: CCNE2

Gene Alias: CYCE2

Gene Summary: The protein encoded by this gene belongs to the highly conserved cyclin family, whose

members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2. This cyclin has been shown to specifically interact with CIP/KIP family of CDK inhibitors, and plays a role in cell cycle G1/S transition. The expression of this gene peaks at the G1-S phase and exhibits a pattern of tissue specificity distinct from that of cyclin E1. A significantly increased expression level of this gene was observed in tumor-derived cells. [provided by RefSeq]