



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

PTK2 recombinant monoclonal antibody

Catalog Number: RAB02694

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against PTK2.

Immunogen: Original antibody is raised against recombinant PTK2.

Theoretical MW (kDa): 125

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: This antibody detects endogenous levels of FAK and does not cross-react with related proteins.

Form: Liquid

Purification: Protein A purification

Isotype: IgG

Recommend Usage: Flow Cytometry (1:50-1:100)

Immunocytochemistry (1:50-1:200)

Immunofluorescence (1:50-1:200)

Immunohistochemistry (1:50-1:200)

Western Blot (1:1000-1:2000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH7.2 (50% glycerol and 0.02% sodium azide)

Storage Instruction: Store at 4°C short term.

Aliquot and store at -20°C long term.

Avoid freeze-thaw cycles.

Entrez GeneID: 5747

Gene Symbol: PTK2

Gene Alias: FADK, FAK, FAK1, pp125FAK

Gene Summary: This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of this gene may be an important early step in cell growth and intracellular signal transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix. At least four transcript variants encoding four different isoforms have been found for this gene, but the full-length natures of only two of them have been determined. [provided by RefSeq]