



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

MMP14 recombinant monoclonal antibody

Catalog Number: RAB02730

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against MMP14.

Immunogen: Original antibody is raised against recombinant MMP14.

Theoretical MW (kDa): 65

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: This antibody detects endogenous levels of MMP14 and does not cross-react with related proteins.

Form: Liquid

Purification: Protein A purification

Isotype: IgG

Recommend Usage: Flow Cytometry (1:50-1:100)

Immunocytochemistry (1:50-1:200)

Immunofluorescence (1:50-1:200)

Immunohistochemistry (1:50-1:200)

Western Blot (1:1000)

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH7.2 (50% glycerol and 0.02% sodium azide)

Storage Instruction: Store at 4°C short term.

Aliquot and store at -20°C long term.

Avoid freeze-thaw cycles.

Entrez GeneID: 4323

Gene Symbol: MMP14

Gene Alias: MMP-X1, MT1-MMP, MTMMP1

Gene Summary: Proteins of the matrix

metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. However, the protein encoded by this gene is a member of the membrane-type MMP (MT-MMP) subfamily; each member of this subfamily contains a potential transmembrane domain suggesting that these proteins are expressed at the cell surface rather than secreted. This protein activates MMP2 protein, and this activity may be involved in tumor invasion. [provided by RefSeq]