



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

CD3E recombinant monoclonal antibody

Catalog Number: RAB02736

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against CD3E.

Immunogen: Original antibody is raised against recombinant CD3E.

Theoretical MW (kDa): 21

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: This antibody detects endogenous levels of CD3-epsilon and does not cross-react with related proteins.

Form: Liquid

Purification: Protein A purification

Isotype: IgG

Recommend Usage: Flow Cytometry (1:50-1:100)
Immunohistochemistry (1:50-1:200)
Western Blot (1:1000-1:2000)
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH7.2 (50% glycerol and 0.02% sodium azide)

Storage Instruction: Store at 4°C short term.
Aliquot and store at -20°C long term.
Avoid freeze-thaw cycles.

Entrez GeneID: 916

Gene Symbol: CD3E

Gene Alias: FLJ18683, T3E, TCRE

Gene Summary: The protein encoded by this gene is the CD3-epsilon polypeptide, which together with CD3-gamma, -delta and -zeta, and the T-cell receptor alpha/beta and gamma/delta heterodimers, forms the T-cell receptor-CD3 complex. This complex plays an important role in coupling antigen recognition to several intracellular signal-transduction pathways. The genes encoding the epsilon, gamma and delta polypeptides are located in the same cluster on chromosome 11. The epsilon polypeptide plays an essential role in T-cell development. Defects in this gene cause immunodeficiency. This gene has also been linked to a susceptibility to type I diabetes in women. [provided by RefSeq]