



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

KIF23 recombinant monoclonal antibody

Catalog Number: RAB02740

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against KIF23.

Immunogen: Original antibody is raised against recombinant KIF23.

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Specificity: This antibody detects endogenous levels of MKLP1 and does not cross-react with related proteins.

Form: Liquid

Purification: Protein A purification

Isotype: IgG

Recommend Usage: Immunocytochemistry
(1:50-1:200)
Immunofluorescence (1:50-1:200)
Western Blot (1:1000-1:5000)
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS, pH7.2 (50% glycerol and 0.02% sodium azide)

Storage Instruction: Store at 4°C short term.
Aliquot and store at -20°C long term.
Avoid freeze-thaw cycles.

Entrez GeneID: 9493

Gene Symbol: KIF23

Gene Alias: CHO1, KNSL5, MKLP-1, MKLP1

Gene Summary: The protein encoded by this gene is a member of kinesin-like protein family. This family

includes microtubule-dependent molecular motors that transport organelles within cells and move chromosomes during cell division. This protein has been shown to cross-bridge antiparallel microtubules and drive microtubule movement in vitro. Alternate splicing of this gene results in two transcript variants encoding two different isoforms. [provided by RefSeq]