



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### PLCG1 recombinant monoclonal antibody, clone PLCG1Y783-C4

**Catalog Number:** RAB02812

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human PLCG1.

**Clone Name:** PLCG1Y783-C4

**Immunogen:** A synthetic phospho-peptide corresponding to residues surrounding Tyr783 of human phospho PLCg1.

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Protein A+G

**Isotype:** Rabbit IgG1k

**Recommend Usage:** Flow Cytometry  
The optimal working dilution should be determined by the end user.

**Storage Buffer:** 1X PBS, 0.02% Sodium azide, 50% Glycerol, 0.1% BSA

**Storage Instruction:** Store at -20°C.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 5335

**Gene Symbol:** PLCG1

**Gene Alias:** PLC-II, PLC1, PLC148, PLCgamma1

**Gene Summary:** The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase

activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]