



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

RELA recombinant monoclonal antibody, clone NFkBS536-C8

Catalog Number: RAB02837

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human RELA.

Clone Name: NFkBS536-C8

Immunogen: A synthetic phospho-peptide corresponding to residues surrounding Ser536 of human phospho NFkB.

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Purification: Protein A+G

Isotype: Rabbit IgG1k

Recommend Usage: Western Blot
The optimal working dilution should be determined by the end user.

Storage Buffer: 1X PBS, 0.02% Sodium azide, 50% Glycerol, 0.1% BSA

Storage Instruction: Store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 5970

Gene Symbol: RELA

Gene Alias: MGC131774, NFkB3, p65

Gene Summary: NFkB1 (MIM 164011) or NFkB2 (MIM 164012) is bound to REL (MIM 164910), RELA, or RELB (MIM 604758) to form the NFkB complex. The p50 (NFkB1)/p65 (RELA) heterodimer is the most abundant form of NFkB. The NFkB complex is inhibited by I-kappa-B proteins (NFkBIA, MIM 164008 or NFkBIB,

MIM 604495), which inactivate NFkB by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, MIM 600664, or IKBKB, MIM 603258) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NFkB complex. Activated NFkB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime (where H is A, C, or T; R is an A or G purine; and Y is a C or T pyrimidine).[supplied by OMIM]