



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### **BTK recombinant monoclonal antibody, clone BtkY551-G12**

**Catalog Number:** RAB02843

**Regulatory Status:** For research use only (RUO)

**Product Description:** Rabbit recombinant monoclonal antibody raised against human BTK.

**Clone Name:** BtkY551-G12

**Immunogen:** A synthetic phospho-peptide corresponding to residues surrounding Tyr551 of human phospho Btk

**Antibody Species:** Rabbit

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Purification:** Protein A+G

**Isotype:** Rabbit IgG1k

**Recommend Usage:** ELISA

Flow Cytometry

Western Blot

The optimal working dilution should be determined by the end user.

**Storage Buffer:** 1X PBS, 0.02% Sodium azide, 50% Glycerol, 0.1% BSA

**Storage Instruction:** Store at -20°C.  
Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 695

**Gene Symbol:** BTK

**Gene Alias:** AGMX1, AT, ATK, BPK, IMD1, MGC126261, MGC126262, PSCTK1, XLA

**Gene Summary:** The protein encoded by this gene plays a crucial role in B-cell development. Mutations in this gene cause X-linked agammaglobulinemia type 1,

which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement. [provided by RefSeq]