



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

MAP2K7 recombinant monoclonal antibody, clone MKK7S271T275-R4F9 (PE)

Catalog Number: RAB02954

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human MAP2K7.

Clone Name: MKK7S271T275-R4F9

Immunogen: A synthetic phospho-peptide corresponding to residues surrounding Ser271 and Thr275 of human MKK7

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Conjugation: PE

Purification: Protein A purification, Protein G purification

Isotype: IgG

Recommend Usage: Flow Cytometry
The optimal working dilution should be determined by the end user.

Storage Buffer: 1X PBS, 0.09% Sodium azide, 0.2% BSA

Storage Instruction: Store at 4°C. Do not freeze.

Entrez GeneID: 5609

Gene Symbol: MAP2K7

Gene Alias: Jnkk2, MAPKK7, MKK7, PRKMK7

Gene Summary: The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase specifically activates

MAPK8/JNK1 and MAPK9/JNK2, and this kinase itself is phosphorylated and activated by MAP kinase kinase kinases including MAP3K1/MEKK1, MAP3K2/MEKK2, MAP3K3/MEKK5, and MAP4K2/GCK. This kinase is involved in the signal transduction mediating the cell responses to proinflammatory cytokines, and environmental stresses. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found, but only one transcript variant has been supported and defined. [provided by RefSeq]