



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

ITGAL recombinant monoclonal antibody, clone MHM24

Catalog Number: RAB03154

Regulatory Status: For research use only (RUO)

Product Description: Mouse recombinant monoclonal antibody raised against human ITGAL.

Clone Name: MHM24

Immunogen: Original antibody is raised against EBV-transformed human lymphoblastoid cells.

Antibody Species: Mouse

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Isotype: IgG1 kappa

Recommend Usage: Flow Cytometry
Inhibition Assay
Immunofluorescence
The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS with 0.02% Proclin 300

Storage Instruction: Store at 4°C for up to 3 months.
For longer storage, aliquot and store at -20°C.
Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3683

Gene Symbol: ITGAL

Gene Alias: CD11A, LFA-1, LFA1A

Gene Summary: ITGAL encodes the integrin alpha L chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form the integrin lymphocyte function-associated antigen-1 (LFA-1), which is expressed on all leukocytes. LFA-1 plays a central role

in leukocyte intercellular adhesion through interactions with its ligands, ICAMs 1-3 (intercellular adhesion molecules 1 through 3), and also functions in lymphocyte costimulatory signaling. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq]