



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

ITGB2 recombinant monoclonal antibody, clone 1B4

Catalog Number: RAB03177

Regulatory Status: For research use only (RUO)

Product Description: Rabbit recombinant monoclonal antibody raised against human ITGB2.

Clone Name: 1B4

Immunogen: Original antibody is raised against monocytes.

Antibody Species: Rabbit

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Isotype: IgG

Recommend Usage: ELISA

Flow Cytometry

Immunofluorescence

Immunoprecipitation

Radioimmunoassay

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS with 0.02% Proclin 300

Storage Instruction: Store at 4°C for up to 3 months.

For longer storage, aliquot and store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3689

Gene Symbol: ITGB2

Gene Alias: CD18, LAD, LCAMB, LFA-1, MAC-1, MF17, MF17

Gene Summary: The product of this gene belongs to the integrin beta chain family of proteins. Integrins are integral cell-surface proteins composed of an alpha chain and a beta chain. This gene encodes the integrin

beta chain beta 2. A given chain may combine with multiple partners resulting in different integrins. For example, beta 2 combines with the alpha L chain to form the integrin LFA-1, and combines with the alpha M chain to form the integrin Mac-1. Integrins are known to participate in cell adhesion as well as cell-surface mediated signalling. Defects in this gene are the cause of leukocyte adhesion deficiency type I (LAD1). Two transcript variants encoding the same protein have been identified for this gene. [provided by RefSeq]