



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Datasheet

### CD1A recombinant monoclonal antibody, clone CBT6

**Catalog Number:** RAB03188

**Regulatory Status:** For research use only (RUO)

**Product Description:** Mouse recombinant monoclonal antibody raised against human CD1A.

**Clone Name:** CBT6

**Immunogen:** Original antibody is raised against human CD1A.

**Antibody Species:** Mouse

**Protocols:** See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

**Form:** Liquid

**Isotype:** IgG1 kappa

**Recommend Usage:** ELISA

Flow Cytometry

Immunohistochemistry

Immunoprecipitation

The optimal working dilution should be determined by the end user.

**Storage Buffer:** In PBS with 0.02% Proclin 300

**Storage Instruction:** Store at 4°C for up to 3 months.

For longer storage, aliquot and store at -20°C.

Aliquot to avoid repeated freezing and thawing.

**Entrez GeneID:** 909

**Gene Symbol:** CD1A

**Gene Alias:** CD1, FCB6, HTA1, R4, T6

**Gene Summary:** This gene encodes a member of the CD1 family of transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of

self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. The protein encoded by this gene localizes to the plasma membrane and to recycling vesicles of the early endocytic system. Alternatively spliced transcript variants have been observed, but their biological validity has not been determined. [provided by RefSeq]