



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Datasheet

KRT7 recombinant monoclonal antibody, clone OV-TL 12/30

Catalog Number: RAB03399

Regulatory Status: For research use only (RUO)

Product Description: Mouse recombinant monoclonal antibody raised against human KRT7.

Clone Name: OV-TL 12/30

Immunogen: Original antibody is raised against ovarian carcinoma cell line OTN11.

Antibody Species: Mouse

Protocols: See our web site at <http://www.abnova.com/support/protocols.asp> or product page for detailed protocols

Form: Liquid

Isotype: IgG1 kappa

Recommend Usage: Flow Cytometry

Immunofluorescence

Immunohistochemistry

Western Blot

The optimal working dilution should be determined by the end user.

Storage Buffer: In PBS with 0.02% Proclin 300

Storage Instruction: Store at 4°C for up to 3 months.

For longer storage, aliquot and store at -20°C.

Aliquot to avoid repeated freezing and thawing.

Entrez GeneID: 3855

Gene Symbol: KRT7

Gene Alias: CK7, K2C7, K7, MGC129731, MGC3625, SCL

Gene Summary: The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and

stratified epithelial tissues. This type II cytokeratin is specifically expressed in the simple epithelia lining the cavities of the internal organs and in the gland ducts and blood vessels. The genes encoding the type II cytokeratins are clustered in a region of chromosome 12q12-q13. Alternative splicing may result in several transcript variants; however, not all variants have been fully described. [provided by RefSeq]